PHILIPPINE JOURNAL OF PSYCHOLOGY, 2015, 48(2), 81-104
COPYRIGHT @ 2015 PSYCHOLOGICAL ASSOCIATION OF THE PHILIPPINES

The Application and Adaptation of Psychological First Aid: The Filipino Psychologists' Experience After Typhoon Haiyan

Bernice Vania N. Landoy
University of the Philippines Visayas

Ma. Regina M. Hechanova
Pia Anna Perfecto Ramos
Ateneo de Manila University

Niel Steve M. Kintanar
University of San Carlos

This study examined the experiences, adaptations, observations, and insights on the use of Psychological First Aid (PFA) in the Philippines. Nineteen PFA providers who were trained in the basics of PFA from three major cities in the Philippines took part in this study. Respondents' knowledge about PFA was consistent with the core principles of promoting a sense of safety, calm, self- and community e cacy, connectedness, and instilling a sense of hope. The delivery of PFA was adapted to the Filipino culture through the use of local counseling models, a group-based set-up, and the use of mindfulness and relaxation techniques. The

Correspondence concerning this article can be addressed to Bernice Vania N. Landoy. Email: bnlandoy@up.edu.ph

Support (MHPSS) in Emergency Settings and began conducting training on the use of PFA. It also exhorted psychologists not to go into areas without assessment and local partners and stressed that psychosocial interventions should not be stand-alone or band-aid solutions but should instead be embedded in integrated and long-term solutions (Hechanova, 2013).

Beyond the principles under PFA, the IASC guidelines on MHPSS in Emergency Settings also recommended the need to utilize culturally-sensitive and evidence-based interventions (IASC, 2007). Litz (2008) described PFA as a þexible process and advocated the need for PFA to take into account the survivorsû culture, ethnicity, religious a ̨ liation, race, and di ̃ ering languages. The WHO, World Trauma Foundation, and World Vision International (2011) guide to PFA also explains that in a crisis situation, a person's spiritual or religious beliefs and rituals may be very important in helping them overcome pain and su ̃ ering, provide meaning to their experience, and give a sense of hope and comfort.

Although PFA is typically done individually, Johnstone (2007) argued that it can be done in groups. Groups allow survivors to share their experiences thereby normalizing their reactions to a stressful situation (Johnstone, 2007). Group sharing also allows survivors to give and receive mutual social support. In fact, the delivery of psychosocial interventions in a group setting has been found to have immediate positive impact and reduce the degree of PTSD symptoms in the long run (Foy et al., 2000).

Everly, Phillips, Kane, and Feldman (2006) outlined the practice of group-PFA. Group-PFA consists of three phases: pregroup activities, six stages of group-PFA, and postgroup activities. The goal of pregroup activities is to perform an initial assessment if it is suitable to provide a group intervention and to check what basic needs are lacking. The six stages of group-PFA include (a) introduction, (b) review or brieýng of what happened, (c) clariýcation or correction about what happened, (d) teaching by o ̃ ering psycho-educational information stress reactions, trauma, and coping, (e) supporting the natural cohesion and resiliency of the group, and (f) assisting in connecting with informal and formal support systems. The goals of postgroup activities include establishing availability for future support and identifying future

needs of higher level of care, volunteers' closure and self-care, and evaluation.

In the Philippines, the use of group-based PFA among government worker-survivors has been found to signiýcantly increase participantsû pre- and posttest scores on self-e cacy and coping skills (Hechanova, Ramos, & Waelde, 2015). The PFA process, psychoeducation and practice of mindfulness, and the use of group sharing were the most useful components of the intervention. Participants cited that group sharing allowed them to share their stories with others who went through the same experience.

The preceding study focused on the beneýts of PFA to survivors. However, the utility of PFA for disaster survivors from the perspective of Filipino providers has not been recorded thus far. Therefore, this study seeks to contribute to the literature by providing information on the knowledge, experience, adaptations made by providers of PFA, and their observations of survivorsû reactions. Speciýcally, we asked:

1) What were the providersû knowledge of PFA and how did responders receive this knowledge?
2) What were their experiences in the conduct of PFA?
3) What adaptations/additions were made in the conduct of the PFA and why?
4) What were survivorsû reactions to PFA?

METHOD

This qualitative study involved interviews/open ended surveys conducted with providers of PFA.

Respondents

Using convenience sampling, 19 PFA providers (14 females and 5 males) participated in this study. Providers reported having undergone basic training on PFA given either by representatives of the World Health Organization, by faculty members of the University of Santo Tomas, or by members of the Psychological Association of the Philippines. Eight came from Metro Manila, six from Cebu, and ýve from Iloilo. Respondersû age ranged from 22 to 51 years old.

might need more advanced support and helping them obtain access to information (WHO et al., 2011).

Process of PFA. When the respondents were asked about the process of PFA, their responses highlighted its dif erence from other forms of intervention like professional counseling, psychological

response teams, and faith-based organizations, and other disaster relief organizations can provide PFA as long as they are trained with the principles and proper delivery. One advantage of PFA compared to

showed that PFA providersû knowledge, conýdence, and readiness increase after training.

Also, according to respondents, PFA is brief and practical. A respondent said, “PFA is brief and it immediately responds to the needs of survivors; more practical.” Another respondent added, “compared to other interventions, [PFA is] not time consuming and does not need further sessions.ò As suggested by the WHO, PFA is often a one-time intervention and providers may only be there to help for a short time (WHO et al., 2011).

Additionally, PFA is easy to follow and use. One respondent said, ñ[PFA is] easy enough, [I] can grasp each of the steps readily,

teaches ways of helping make people's burden easier to bear. In this way, the *pagdadala* model is also considered a community-based mental health and psychosocial support model (M. L. Verzosa, personal communication, December 8, 2015). Two PFA providers from Iloilo[1], for instance, said:

> The pagdadala model of Decenteceo, where respondents expressed their needs and the burdens they are carrying in life with a group and explored or presented ways and means in dealing with their concerns. Respondents were not categorically directed to talk about their experiences during the disaster. Some

was to run PFA in groups. This was done not only because of the lack of facilitators but also because it is believed that Filipinos enjoy being in groups. A local study showed that Filipino participants in group-based PFA found sharing their experiences with others a useful component of the intervention because it made them feel a sense of solidarity with other survivors who had undergone the same experience (Hechanova et al., 2015) As one respondent said, ñthe spirit of PFA is, if you really think about it, is *bayanihan*[2]."

Filipino PFA providers were able to follow the dĩ erent phases of group-PFA (Everly et al., 2006). Pregroup activities were conducted during meetings with the psychosocial support organizers and a needs assessment survey was distributed while waiting for participants to arrive. Asking and setting of expectations were usually the beginning of conducting group-PFA. During *kamustahan*, participants were usually divided into small groups of six to 10 members but during the psychoeducation stage, participants were gathered in one big group. Also, as part of postgroup activities, PFA included discussions on future care and the support these survivors need.

Open-space activity and PFA. The PFA providers also mentioned that they use open-space activity when they see that participants are not ready to individually talk about their concerns. In particular, one PFA provider said:

> We used an open-space activity where individuals go around and write solutions to the problems they identiýed on manila papers posted on the walls. A person (or administrator) was there to provide answers. They had so many basic needs that were unmet so the open space activity along with linkages to authority was crucial.

Challenges in Conducting PFA

However, respondents also cited the challenges in the delivery of PFA and these were: (a) lack of protocols for serious cases and for follow up, (b) lack of conducive venues, (c) linking with services, (d) lack of disaster response systems, and (e) the limited number of trained individuals who can deliver PFA.

[2]Bayanihan is a Filipino term which means the spirit of communal unity.

Respondents also recognized that PFA might not be su cient for people who have been identi ed or diagnosed with mental health disorders or are severely distressed, which is in keeping with international guidelines (Cain et al., 2010). As mentioned, the respondents cited the lack of protocol on what to do for people who need more specialized help as another challenge. They acknowledged that other than providing PFA, they also needed to refer survivors, particularly those who are vulnerable or at-risk, to other services. Despite the need for these, there were no mechanisms for referral, documentation, or follow up.

The respondents also cited the lack of conducive venues as another challenge in the delivery of PFA in the Philippines. They reported di culty in facilitating relaxation techniques in some areas because of the noise.

Aside from these, the respondents acknowledged that PFA needs to be delivered with partners or a network, otherwise there will not be any follow through. They also emphasized the value of linking people with practical support and social support like family and the appropriate agencies, which WHO describes as a major part of PFA (WHO et al., 2011). However, they reported di culty in linking the survivors with agencies due to "poor networking in government agencies." The respondents also said that the lack of a systematic disaster response system in the Philippines limits the support that they can provide to the survivors. One PFA provider said, "you cannot give practical assistance yourself because you are there to give social support and other kinds of help. How can you be running around?"

Finally, because there was only a small number of people who could deliver PFA, respondents had to shift to group-based PFA because of the need to assist as many survivors as possible.

Perceived Survivors' Reactions to PFA

When respondents were asked what reactions they observed among the survivors after conducting PFA, their answers can be summarized into three: physical and behavioral, cognitive, and emotional.

Responders reported physical and behavioral reactions such as

a change in facial expression, smiling, and expressing their gratitude among the survivors after the PFA session. One PFA provider shared that:

> At ýrst you can really see how burdened they are through their facial expression and then after, they feel relieved, their faces light up. Some are hesitant to trust you at ýrst and, then in the end, their smiles become genuine and they are thankful.

According to the respondents, survivors were smiling after the session and were expressing their gratitude immensely. ñThey [said] thank you for listening, for giving them a chance to express especially since itûs within a group settingò and ñThankful [that] they got someone who cared for them and that their needs and concerns were heard and that they got to become more positive."

Another observation was the gradual cognitive change among the survivors. In particular, the recipients were able to reframe the way they see themselves, from being victims to survivors. They also understood that their stress reactions were because of the event, and recognized that these reactions are normal. According to the respondents, the survivors acknowledged that they were not prepared for the calamity but at the same time believed that they will be okay. A respondent shared, ñAt ýrst, the survivors were confused because they were not aware what PFA was all about, and then, through the psychoeducation part, they began to understand and appreciate it." Another respondent shared the insights of some survivors saying, "They never realized that they were so stressed because they kept on working. They thought they were okayò and ñ[they] acknowledge that, although not all needs will be met just yet, but on a psychological level, they have this, 'mas kaya ko na ngayon' (I can do this) mentality."

Responders also reported changes in emotions, such as emotions of relief, gratitude, and hope, among survivors as a result of PFA. The respondents shared that the survivors felt relieved from the stress and they felt the burden was lessened. One respondent said: "From negative feelings, having trauma and þashbacks, to already having hope when you begin to link them and when they already know who they can approach to ask for help."

The respondents attributed this feeling of hope among the survivors to the knowledge that help is available when they need it and

the realization that there are people who will care for them.

Limitations and Implications for Future Research

As this study is exploratory in nature, there remains a great need for further ýeld research to evaluate the delivery and e ectiveness of PFA in a variety of postdisaster contexts here in the Philippines. For one, the study focused only on the providers of PFA who were trained in such intervention and might have a favorable response bias. Hence, research that looks at PFA from the point of view of survivors is also important.

It is also very relevant to note that the results reþect the perceived utility of PFA from PFA providers' perspectives as opposed to a more objective measure of the actual utility of PFA in the ýeld. This could be addressed by conducting more research on PFAîs e ectiveness in a number of contexts and methods (Forbes et al., 2011). An example of this is a ýeld experiment comparing survivors who received PFA and those who did not. This is imperative to establish the impact of PFA.

Another limitation of the study is that majority of the respondents of the study have a background in psychology and/or received postgraduate training in counseling/clinical psychology. Comparison studies between perception of professionals and non-professionals on the implementation of PFA can also be explored to see whether the reactions and insights on the use of PFA will be the same.

Implications and Conclusion

Despite its limitations, the results of this exploratory study show that over and above the additions and adaptations done, the knowledge of PFA among Filipino psychologists is consistent with its purpose and nature as outlined by current guidelines and literature (Brymer, 2006; WHO et al., 2011). The providers were able to retain what they have learned in the trainings that they received and applied these in the ýeld. Providers of PFA were conýdent of its value and were satisýed with their experience of conducting PFA because of the reactions and changes of the survivors that they observed and because of its ease of use as opposed to the commonly practiced CISD. Although,

additions and adaptations were made to the steps, the nature of PFA's þexibility ð individual versus group, timing, context, preference of survivor, and culture—is its strength. This is line with the assertion of

Perspectives in Psychiatric Care, 43, 38-40.
Kantor, E. M., & Beckert, D. R. (2011). Psychological ýrst aid. In F. J. Stoddard, A. A. Pandya, & C. L. Katz (Eds.),

psychological intervention since 9/11. *American Psychologist, 66*(6), 482-492. doi:10.1037/a0024806

WHO-Assessment Instruments for Mental Health Systems (2006).